

The Born and Lens-Lens Corrections to Weak Gravitational Lensing Angular Power Spectra

Charles Shapiro

Department of Physics, The University of Chicago, Chicago, IL 60637
Kavli Institute for Cosmological Physics, The University of Chicago, Chicago, IL 60637

Asantha Cooray

Center for Cosmology, Department of Physics and Astronomy, University of California, Irvine, CA 92617

Abstract. We revisit the estimation of higher order corrections to the angular power spectra of weak gravitational lensing. Extending a previous calculation of Cooray and Hu, we find two additional terms to the fourth order in potential perturbations of large-scale structure corresponding to corrections associated with the Born approximation and the neglect of line-of-sight coupling of two foreground lenses in the standard first order result. These terms alter the convergence ($\kappa\kappa$), the lensing shear E-mode ($\epsilon\epsilon$), and their cross-correlation ($\kappa\epsilon$) power spectra on large angular scales, but leave the power spectra of the lensing shear B-mode ($\beta\beta$) and rotational ($\omega\omega$) component unchanged as compared to previous estimates. The new terms complete the calculation of corrections to weak lensing angular power spectra associated with both the Born approximation and the lens-lens coupling to an order in which the contributions are most significant. Taking these features together, we find that these corrections are unimportant for any weak lensing survey, including for a full sky survey limited by cosmic variance.

1. Introduction

Weak gravitational lensing provides a powerful way to probe the matter distribution of the large-scale structure of the Universe and to measure cosmological parameters both through growth of structures and geometrical distance projections (see, [1, 2, 3] for recent reviews). As weak lensing experiments improve in precision, it is necessary to improve the accuracy to which lensing statistics are computed. Included in the sources of theoretical uncertainties are various approximations used in the canonical calculation of the weak lensing power spectra. Two such approximations, the Born approximation and the exclusion of lens-lens coupling involving two foreground lenses along the line-of-sight, were investigated by Cooray and Hu [4] using analytical methods, while numerical simulations have also been used to explore these approximations [10, 12]. Relaxing these approximations introduce sub-percent-level corrections to the power spectra of the convergence (κ) and the shear E-mode (ϵ). Miniscule contributions to the angular power spectra of shear B-mode (β) and image rotational (ω) components are also generated, though in the standard calculation, these two components are exactly zero. This has led to the use of the shear B-mode component monitoring systematics in lensing studies.

In Cooray and Hu [4], the estimation of lensing corrections was done perturbatively; they computed corrections to the power spectra which were fourth order in the gravitational potential of large-scale structure. In this paper, we briefly review these corrections and show that there exist two additional terms of the same order. Mainly, these terms decrease the power of the convergence and shear ϵ -mode on large angular scales. Neither the shear β -mode nor the rotational component are changed relative to the previous estimate in Ref. [4]. Together, the calculation is complete and we can make a strong conclusion that the corrections due to the Born approximation and lens-lens coupling are negligible for both current and upcoming lensing experiments. Unless one begins to probe multipole moments greater than 10^4 , it is likely that these corrections can be ignored for any experiment, including an ultimate all-sky experiment limited by cosmic variance alone. While the calculation related to higher order corrections is complete, there may be other sources of error in the theoretical estimates of lensing statistics [2, 5]. For example, theoretical estimates of the underlying non-linear dark matter clustering is not known to better than 10% at arcminute scales [6] and is likely to be further complicated by baryon physics [13, 14]. Another source of important error is that lensing statistics are based on reduced shear rather than shear directly, which results in perturbative corrections from convergence to shear [15]. Clustering of background sources also contaminates weak lensing statistics [7] and may be an important source of uncertainty for tomographic lensing measurements [9] in which the background source redshift distribution is binned to obtain lensing measurements as a function of redshift. Based on our calculation, we are now confident that these remaining issues dominate the theoretical calculation and interpretation of the weak lensing angular power spectra.

This brief paper is organized as following: In the next Section, we outline the

calculation related to higher order corrections to weak lensing angular power spectrum. We identify two new terms by extending the calculation of Ref. [4], and make numerical estimates of the correction. We find that the overall correction is negligible even for an all-sky lensing experiment limited by cosmic variance and probing fluctuations out to a multipole of 10^4 in the angular power spectrum. We conclude with a brief summary of our results in Section 3. For illustrative purposes, we assume a flat Λ CDM cosmological model with parameters $\Omega_c = 0.3$, $\Omega_b = 0.05$, $\Omega_\Lambda = 0.65$ for CDM, baryon, and cosmological constant densities relative to the closure density, a scaled Hubble constant of $h = 0.7$, and a primordial power spectrum of fluctuations normalized to $\sigma_8 = 0.85$ with a tilt of $n = 1$.

2. Calculational Method

Here we provide a brief summary of the analytical calculation. For further details, we refer the reader to Cooray & Hu [4]. For simplicity, we also use their notation throughout.

The distortion tensor for a weakly lensed source is given by an integral over a photon's path, $\mathbf{x}$, through the large-scale gravitational potential:

$$\psi_{ab}(\mathbf{n}, \chi_s) = 2 \int_0^{\chi_s} d\chi g(\chi, \chi_s) \Phi_{,ac}(\mathbf{x}; \chi) [\delta_{cb} + \psi_{cb}(\mathbf{n}, \chi)], \quad (1)$$

where $\mathbf{n}$ is the source's sky position, χ_s is its comoving distance from us, Φ is the potential and δ is the Kronecker delta. Commas denote spatial derivatives in the transverse directions, and there is an implicit sum over repeated indices. The weighting function g assumes that the source is at a fixed distance away; it is given by

$$g(\chi', \chi) \equiv \begin{cases} d_A(\chi - \chi') d_A(\chi') / d_A(\chi) & \text{for } \chi' < \chi \\ 0 & \text{for } \chi' \geq \chi \end{cases} \quad (2)$$

with d_A as the angular diameter distance. The power spectra of weak lensing observables are found by combining components of C_{abcd} , defined as

$$\langle \psi_{ab}^*(\mathbf{l}) \psi_{cd}(\mathbf{l}') \rangle \equiv (2\pi)^2 \delta(\mathbf{l} - \mathbf{l}') C_{abcd}(\mathbf{l}), \quad (3)$$

with $\psi(\mathbf{l})$ the 2D Fourier transform of $\psi(\mathbf{n})$.

The canonical calculation of the power spectra takes advantage of the fact that the potential – and therefore the deflection of light – is small. Lens-lens coupling, i.e. the appearance of ψ on the right side of equation Eq. 1, can clearly be ignored to first order in Φ . Furthermore, since the photon path is nearly a straight line, we may write

$$\mathbf{x}(\mathbf{n}, \chi) = \mathbf{n} d_A(\chi) + \delta \mathbf{x}(\mathbf{n}, \chi), \quad (4)$$

and in the Born approximation set the transverse deflection $\delta \mathbf{x}$ to zero. The result is a simplified integral over the undeflected path,

$$\psi_{ab}(\mathbf{n}, \chi_s) = 2 \int_0^{\chi_s} d\chi g(\chi, \chi_s) \Phi_{,ab}(\mathbf{n} d_A(\chi); \chi) + O(\Phi^2), \quad (5)$$

which, in the Limber approximation [16], leads to the standard expression for the two point functions of ψ in terms of the power spectrum of Φ [17]

$$C_{abcd}(\mathbf{l}) = 4l_a l_b l_c l_d \int d\chi \frac{g(\chi, \chi_S)^2}{d_A^6(\chi)} P(l/d_A; \chi). \quad (6)$$

The first order spectra of the lensing observables $(\kappa, \epsilon, \beta, \omega)$ are

$$C_l^{\kappa\kappa} = C_l^{\epsilon\epsilon} = C_l^{\beta\beta} = l^4 \int d\chi \frac{g(\chi, \chi_S)^2}{d_A^6(\chi)} P(l/d_A; \chi) \quad (7)$$

with all other combinations equal to zero. Note that throughout this paper, we will be employing Limber approximation to calculate angular power spectra. This assumption makes use of the flat-sky approximation and could potentially affect the large angular scale correlations. However, as a test case, we considered the calculation exactly by integrating over spherical Bessel functions that one encounters with the all-sky projections and found our results to be accurate at the percent level and below at $\ell > 10$. Just as the Limber approximation is adequate to calculate convergence power spectrum at the first order (e.g., Ref. [11] based on simulations), we believe it is adequate for the calculations at the higher order.

2.1. Perturbative Corrections

To relax the Born approximation, we simply Taylor expand the potential in equation Eq. 1 about the undeflected photon path. With the transverse deflection given by

$$\delta x_a(\mathbf{n}, \chi) = -2 \int d\chi' g(\chi', \chi) \frac{d_A(\chi)}{d_A(\chi')} \Phi_{,a}(\mathbf{x}; \chi'), \quad (8)$$

the potential is

$$\begin{aligned} \Phi(\mathbf{x}; \chi) &= \Phi(\mathbf{n}d_A(\chi) + \delta\mathbf{x}; \chi) = \Phi(\mathbf{n}d_A; \chi) + \delta x_a \Phi_{,a}(\mathbf{n}d_A; \chi) \\ &\quad + \frac{1}{2} \delta x_a \delta x_b \Phi_{,ab}(\mathbf{n}d_A; \chi) + O(\Phi^4). \end{aligned} \quad (9)$$

Note that $\delta\mathbf{x}$ depends on the deflected path, and must also be expanded perturbatively. Substituting equation Eq. 9 into equation Eq. 8, we find the deflection to second order in potential fluctuations to be:

$$\delta x_a(\mathbf{n}, \chi) = \delta x_a^{(1)}(\mathbf{n}, \chi) + \delta x_a^{(2)}(\mathbf{n}, \chi) + O(\Phi^3) \quad (10)$$

$$\delta x_a^{(1)}(\mathbf{n}, \chi) \equiv -2 \int d\chi' g(\chi', \chi) \frac{d_A(\chi)}{d_A(\chi')} \Phi_{,a}(\mathbf{n}d_A(\chi'); \chi') \quad (11)$$

$$\delta x_a^{(2)}(\mathbf{n}, \chi) \equiv -2 \int d\chi' g(\chi', \chi) \frac{d_A(\chi)}{d_A(\chi')} \Phi_{,ab}(\mathbf{n}d_A(\chi'); \chi') \delta x_b^{(1)}(\mathbf{n}, \chi'). \quad (12)$$

Now we may express $\Phi(\mathbf{x})$, the potential at the deflected position, using integrals over the undeflected path:

$$\begin{aligned} \Phi(\mathbf{x}; \chi) &= \Phi(\mathbf{n}d_A; \chi) + (\delta x_a^{(1)} + \delta x_a^{(2)}) \Phi_{,a}(\mathbf{n}d_A; \chi) \\ &\quad + \frac{1}{2} \delta x_a^{(1)} \delta x_b^{(1)} \Phi_{,ab}(\mathbf{n}d_A; \chi) + O(\Phi^4). \end{aligned} \quad (13)$$

Inserting equation Eq. 13 into equation Eq. 1 leads to ψ as a line-of-sight projection of a source field,

$$\psi_{ab}(\mathbf{n}, \chi_S) = 2 \int d\chi g(\chi, \chi_S) S_{ab}(\mathbf{n}d_A; \chi), \quad (14)$$

where $S_{ab}(\mathbf{n}d_A; \chi)$ includes the following terms (all depend on χ):

$$S_{ab}^{(1)} = \Phi_{,ab}(\mathbf{n}d_A) \quad (15)$$

$$S_{ab}^{(2B)} = \delta x_c^{(1)}(\mathbf{n}) \Phi_{,abc}(\mathbf{n}d_A) \quad (16)$$

$$S_{ab}^{(3X)} = \delta x_c^{(2)}(\mathbf{n}) \Phi_{,abc}(\mathbf{n}d_A) \quad (17)$$

$$S_{ab}^{(3B)} = (1/2) \delta x_c^{(1)}(\mathbf{n}) \delta x_d^{(1)}(\mathbf{n}) \Phi_{,abcd}(\mathbf{n}d_A). \quad (18)$$

These terms yield the standard second order expression for C_{abcd} plus higher order corrections. Third order corrections vanish in the Limber approximation. Cooray and Hu [4] computed all fourth order corrections except for a term that couples $S^{(3X)}$ to $S^{(1)}$. We find that it is given by

$$C_{abcd}^X(\mathbf{l}) = 4 \int d\chi \frac{g(\chi, \chi_S)^2}{d_A(\chi)^6} P_{abcd}^X(\mathbf{l}; \chi) \quad (19)$$

$$P_{abcd}^X(\mathbf{l}; \chi) = -4 \int \frac{d^2 l'}{(2\pi)^2} (l_a l_b l'_c l'_d + l'_a l'_b l_c l_d) (\mathbf{l} \cdot \mathbf{l}')^2 \int d\chi' \frac{g(\chi', \chi)^2 d_A(\chi)^2}{d_A(\chi')^8} \\ \times P\left(\frac{l}{d_A(\chi)}; \chi\right) P\left(\frac{l'}{d_A(\chi')}; \chi'\right), \quad (20)$$

where P_{abcd}^X is the correction to the power spectrum of the source field. The corrections to the power and cross spectra of the lensing observables ($\kappa, \epsilon, \beta, \omega$) are

$$C_l^{X\alpha\beta} = -\frac{\eta}{\pi} l^4 \int d\chi \frac{g(\chi, \chi_S)^2}{d_A(\chi)^4} \int dl' l'^5 \\ \times \int d\chi' \frac{g(\chi', \chi)^2}{d_A(\chi')^8} P\left(\frac{l}{d_A(\chi)}; \chi\right) P\left(\frac{l'}{d_A(\chi')}; \chi'\right), \quad (21)$$

with

$$\eta \equiv \begin{cases} 2 & \text{for } \alpha\beta = \kappa\kappa \\ 3/2 & \text{for } \alpha\beta = \kappa\epsilon \\ 1 & \text{for } \alpha\beta = \epsilon\epsilon \\ 0 & \text{otherwise} \end{cases}. \quad (22)$$

Similarly, lens-lens coupling can be accounted for by iteratively expanding the recursive term in equation Eq. 1. This yields the following second and third order source terms:

$$S_{ab}^{(2L)} = -2\Phi_{,ac}(\mathbf{n}d_A; \chi) \int d\chi' g(\chi', \chi) \Phi_{,cb}(\mathbf{n}d_A(\chi'); \chi') \quad (23)$$

$$S_{ab}^{(3L)} = 4\Phi_{,ac}(\mathbf{n}d_A; \chi) \int d\chi' g(\chi', \chi) \Phi_{,cd}(\mathbf{n}d_A(\chi'); \chi') \\ \times \int d\chi'' g(\chi'', \chi') \Phi_{,ab}(\mathbf{n}d_A(\chi''); \chi''). \quad (24)$$

The first-second order correction to C_{abcd} vanishes under the Limber approximation. Cooray and Hu [4] calculated the second-second order correction, but neglected the

correction involving the first and the third order terms. This term does not vanish; it can be written as

$$P_{abcd}^{(13)} = d_A(\chi)^{-6} \int d\chi' \int d^2n e^{i\mathbf{l}\cdot\mathbf{n}} \langle S_{ab}^{*(1)}(\mathbf{n}d_A(\chi); \chi) S_{cd}^{(3L)}(\mathbf{0}; \chi') \rangle + (ab \leftrightarrow cd) \quad (25)$$

$$= 4d_A(\chi)^{-6} \iiint d\chi' d\chi'' d\chi''' \int d^2n e^{i\mathbf{l}\cdot\mathbf{n}} g(\chi'', \chi') g(\chi''', \chi'') \times \langle \Phi_{,ab}(\mathbf{n}d_A(\chi); \chi) \Phi_{,ce}(\mathbf{0}; \chi') \Phi_{,ef}(\mathbf{0}; \chi'') \Phi_{,fd}(\mathbf{0}; \chi''') \rangle + (ab \leftrightarrow cd). \quad (26)$$

To complete the derivation, each of the four Φ s above must be expanded into its Fourier modes, $\phi(\mathbf{k})$. Then to calculate the ensemble average, $\langle \phi(\mathbf{k})\phi(\mathbf{k}')\phi(\mathbf{k}'')\phi(\mathbf{k}''') \rangle$, one must sum over all possible pairs of Wick contractions of the ϕ s. Contracting two ϕ s ultimately produces delta functions which match up the corresponding χ s. For example, these contractions

$$\overbrace{\langle \phi(\mathbf{k})\phi(\mathbf{k}')\phi(\mathbf{k}'')\phi(\mathbf{k}''') \rangle}$$

lead to a term containing $\delta(\chi - \chi'')\delta(\chi' - \chi''')$. In fact, this is the only non-trivial term: because of the weighting functions in Eq. 26, delta functions in other terms ($\delta(\chi'' - \chi')$ and $\delta(\chi'' - \chi''')$) will cause the integrals to vanish [4]. Nonetheless, the first-third correction does not vanish completely, and we find that $P_{abcd}^{(13)}$ is coincidentally equal to P_{abcd}^X in equation Eq. 20.

There also exist corrections due to couplings between Born and lens-lens source terms. However, in this case, we do not find additional corrections apart from those already considered by Cooray and Hu [4]. To the highest order in importance, i.e., fourth order in potential perturbations, this completes the calculation of corrections to weak lensing angular power spectra.

3. Results & Summary

We have calculated the $O(\Phi^4)$ corrections to weak lensing power spectra that arise from the Born approximation and lens-lens coupling; they are illustrated in Fig. Fig. 1. The primary effect of the new term, C_l^X , is to make the total correction to the $\kappa\kappa$, $\kappa\epsilon$ and $\epsilon\epsilon$ spectra negative for $l \ll 200$. The β and ω spectra are unaffected by the new term when comparing to the calculation of Ref. [4]. As discussed in [4], the ω term arises from lens-lens coupling only; it is not a test of the Born approximation as was suggested in Ref. [10]. The larger difference in the amplitude of the rotational power spectrum shown here and the one shown in Ref. [8] (their Figure 2) is due to the large difference in the source redshift used between these two calculations: here we use $z_s = 1$ for galaxy lensing surveys while $z_s = 1100$ is used in [8] for lensing of cosmic microwave background anisotropies. The large difference suggests that non-linear couplings that lead to these second order corrections grow significantly with increasing path length between source and observer, though the correction still remains below the first order

term. The conclusions of Cooray and Hu are unchanged: corrections related to the Born approximation and lens-lens coupling can be safely neglected for current surveys. They do not limit the use of β -modes in monitoring systematic errors.

Furthermore, since our calculation is now complete, we can make even a stronger statement. As shown in Figure Fig. 1, the higher order corrections are below the cosmic variance level of an all-sky survey, $\sqrt{2/(2l+1)}C_l$, out to a multipole of 10^4 when these corrections start to be higher than the cosmic variance level. This suggests that for all experiments that use statistical information out to a multipole of 10^4 , the angular power spectra will not be affected by the corrections associated with the Born approximation and the coupling of two lenses. This, however, does not imply that theoretical estimates of the angular power spectra are accurate to such a high precision. Throughout this paper, we have relied on the Limber approximation, which ignores contributions to the power spectra from modes with wave-vectors parallel to the line-of-sight. This approximation is justified by lensing simulations [11], which agree with the canonical calculation of the power spectra and hence our conclusions at the few percent level. We have also assumed that the weak lensing calculation is valid, despite the fact that the lensing observables, such as convergence, have non-gaussian distributions with tails that can affect the non-linear part of their spectra, depending on how one includes high lensing peaks that merge weak lensing scales to strong lensing in the data analysis [12]. Beyond issues related to the weak lensing approximation, the non-linear matter power spectrum for a given cosmology is uncertain at the 10% level, both due to uncertainties in the mapping between linear to non-linear dark matter fluctuations and the effect of baryons in determining how dark matter clusters on arcminute scales and below [13, 14]. Accounting for such uncertainties are beyond those possible with analytical calculations. In fact, all remaining uncertainties are those that must be addressed with numerical simulations. We can now be confident that the basic analytical formulation of lensing power spectra is well explored and the first order result is accurate enough for all upcoming experiments.

To summarize, here we have revisited the estimation of higher order corrections to the angular power spectra of weak gravitational lensing and have extended the calculation of Cooray and Hu [4] that first described the corrections related to the Born approximation and the neglect of line-of-sight coupling of two foreground lenses in the standard first order result. We found two additional terms to the fourth order in potential perturbations of the large-scale structure and these terms, in return, altered the convergence ($\kappa\kappa$), the lensing shear E-mode ($\epsilon\epsilon$), and their cross-correlation ($\kappa\epsilon$) power spectra on large angular scales. There is no modification to the lensing shear B-mode ($\beta\beta$) and rotational ($\omega\omega$) component power spectra when compared to the previous estimate. With these new terms, the calculation of corrections to weak lensing power spectrum associated with both the Born approximation and the lens-lens coupling is complete. The overall numerical result is that these corrections are unimportant for any weak lensing survey including for a full sky survey limited by the cosmic variance.




Figure 1. Left: Weak lensing power spectrum and corrections. The solid line shows the first order spectrum; the dot-dashed line shows the new correction, $C_l^{X\epsilon\epsilon}$ ($\eta = 1$), which is negative; the dotted and dashed lines show the full corrections (Born and lens-lens) to the $\kappa\kappa$ and $\epsilon\epsilon$ spectra respectively – they are negative at low l and positive at high l as indicated by (-)/(+) signs. For reference, we also show the cosmic variance level of an all-sky experiment with a thin long-dashed line. The comparison shows that for $l < 10^4$, the full corrections from Born and lens-lens coupling are below the cosmic variance level and is unlikely to be an error of statistical significance. Right: Full corrections divided by the first order result. The dotted and dashed lines show the relative corrections for $\kappa\kappa$ and $\epsilon\epsilon$ respectively, while the thin long-dashed line is the ratio of cosmic variance to the first order result.

4. Acknowledgments

Authors thank Wayne Hu for helpful discussions, Martin White and Chris Vale for comments, and Scott Dodelson for inspiring this paper. CAS was supported in part by the Department of Energy, the Kavli Institute for Cosmological Physics at the University of Chicago and by NSF grant PHY-011442.

References

- [1] M. Bartelmann and P. Schneider, Phys. Rept. **340**, 291 (2001) [arXiv:astro-ph/9912508]; P. Schneider, arXiv:astro-ph/0509252.
- [2] A. Refregier, Ann. Rev. Astron. Astrophys. **41**, 645 (2003) [arXiv:astro-ph/0307212].
- [3] P. Schneider, arXiv:astro-ph/0509252.
- [4] A. Cooray and W. Hu, Astrophys. J. **574**, 19 (2002) [arXiv:astro-ph/0202411].
- [5] B. Jain, M. Jarvis and G. Bernstein, arXiv:astro-ph/0510231.
- [6] R. E. Smith *et al.* [The Virgo Consortium Collaboration], Mon. Not. Roy. Astron. Soc. **341**, 1311 (2003) [arXiv:astro-ph/0207664].
- [7] F. Bernardeau A&A **338**, 375 (1998) [arXiv:astro-ph/9712115]; A. Cooray preprint (2002) [arXiv:astro-ph/0202465].
- [8] C. M. Hirata and U. Seljak, Phys. Rev. D **68**, 083002 (2003) [arXiv:astro-ph/0306354].

- [9] B. Jain & A. Taylor, *Phys. Rev. Lett.* **91**, 141302 (2003) [arXiv:astro-ph/0306046]; M. Takada & M. White, *Astrophys. J.* **601**, L1 (2004) [arXiv:astro-ph/0311104].
- [10] B. Jain, U. Seljak and S. D. M. White, *Astrophys. J.* **530**, 547 (2000) [arXiv:astro-ph/9901191].
- [11] M. J. White and W. Hu, *Astrophys. J.* **537**, 1 (2000) [arXiv:astro-ph/9909165].
- [12] C. Vale and M. J. White, *Astrophys. J.* **592**, 699 (2003) [arXiv:astro-ph/0303555].
- [13] M. J. White, *Astropart. Phys.* **22**, 211 (2004) [arXiv:astro-ph/0405593];
- [14] H. Zhan and L. Knox, *Astrophys. J.* **616**, L75 (2004) [arXiv:astro-ph/0409198].
- [15] M. J. White, *Astropart. Phys.* **23**, 349 (2005) [arXiv:astro-ph/0502003]; S. Dodelson, C. Shapiro and M. J. White, arXiv:astro-ph/0508296.
- [16] D. Limber, *Astrophys. J.* **119**, 655 (1954)
- [17] N. Kaiser, *Astrophys. J.* **388**, 286 (1992)